



Side Effects

A significant hormonal transition will occur in the first four weeks after the insertion of your hormone pellets. Therefore, certain changes might develop that can be bothersome.

FLUID RETENTION: Testosterone stimulates the muscle to grow and retain water, which may result in a weight change of two to five pounds. This is only temporary. This happens frequently with the first insertion, and especially during hot, humid weather conditions.

SWELLING OF THE HANDS & FEET: This is common in hot and humid weather. It may be treated by drinking lots of water, reducing your salt intake, taking cider vinegar capsules daily, (found at most health and food stores) or by taking a mild diuretic, which the office can prescribe.

UTERINE SPOTTING/BLEEDING: This may occur in the first few months after an insertion, especially if you have been prescribed progesterone and are not taking properly: i.e. missing doses, or not taking a high enough dose. Please notify the office if this occurs. Bleeding is not necessarily an indication of a significant uterine problem. More than likely, the uterus may be releasing tissue that needs to be eliminated. This tissue may have already been present in your uterus prior to getting pellets and is being released in response to the increase in hormones.

MOOD SWINGS/IRRITABILITY: These may occur if you were quite deficient in hormones. They will disappear when enough hormones are in your system. 5HTP can be helpful for this temporary symptom and can be purchased at many health food stores.

FACIAL BREAKOUT: Some pimples may arise if the body is very deficient in testosterone. This lasts a short period of time and can be handled with a good face cleansing routine, astringents and toner. If these solutions do not help, a link to purchase a natural supplement called DIM can be sent to you at any time, OR if you prefer a prescription, we can call in spironolactone for you. If you're already prone to this condition, please let us know so you can start either medication now. Spironolactone will require a potassium level check at 6 weeks after starting and every 6 months while you remain on the medication because it CAN increase your potassium above safe levels.

1113 S. Range Avenue, Ste 140 Denham Springs, LA 70726
702 Chestnut Street Murphysboro, IL 62966
(O): 225-283-1211 (F): 225-283-1217



HAIR LOSS: Is rare and usually occurs in patients who convert testosterone to DHT. Dosage adjustment generally reduces or eliminates the problem. Prescription medications may be necessary in rare cases.

HAIR GROWTH: Testosterone may stimulate some growth of hair on your chin, chest, nipples and/or lower abdomen. This tends to be hereditary. You may also have to shave your legs and arms more often. Dosage adjustment generally reduces or eliminates the problem. We offer semi-permanent laser hair growth that will provide a more long-term solution.

1113 S. Range Avenue, Ste 140 Denham Springs, LA 70726
702 Chestnut Street Murphysboro, IL 62966
(O): 225-283-1211 (F): 225-283-1217